

First record of the Mediterranean horseshoe bat *Rhinolophus euryale* (BLASIUS, 1853) in Austria (Chiroptera, Rhinolophidae)

FRIEDERIKE SPITZENBERGER^{1*}, EDMUND WEISS¹ & PETER SACKL²

¹ BatLife Österreich, Landstraßer Hauptstraße 139/15, 1030 Wien, Austria, friederike.spitzenberger@aon.at

² Universalmuseum Joanneum, Studienzentrum Natur, Weinzöttlstr. 16, 8045 Graz-Andritz, Austria

Erstnachweis der Mittelmeer-Hufeisennase *Rhinolophus euryale* (BLASIUS, 1853) in Österreich (Chiroptera, Rhinolophidae)

Abstract

On 6 January 2022, a solitary Mediterranean horseshoe bat (*Rhinolophus euryale*) was observed in the Lurgrotte Cave in Semriach, Styria, Austria (47° 13' N, 15° 22' E). This cave is located at the southern slope of the Eastern Alps at 670 m a. s. l. The species was identified by the noticeable lilac tinge of its fur and size intermediate between greater and lesser horseshoe bat. This sighting constitutes the first record of *Rhinolophus euryale* in Austria.

Key words

Rhinolophus euryale, Austria

On 6 January 2022, the authors performed together with Wolfgang and Mathias Moche (both Vienna) the annual census of bats hibernating in karst caves along the middle reaches of the Mur river in Styria, south east Austria (SACKL et al. 2011). In the Lurgrotte Cave, a solitary medium-sized rhinolophid bat stood out from 1277 lesser (*Rhinolophus hipposideros*) and 32 greater horseshoe bats (*Rh. ferrumequinum*). According to size and colouration we identified it as Mediterranean horseshoe bat (*Rhinolophus euryale*).

Zusammenfassung

Am 6. Januar 2022 beobachteten wir eine einzelne Mittelmeer-Hufeisennase in der Lurgrotte in Semriach, Steiermark, Österreich (47° 13' N, 15° 22' E). Diese Höhle liegt am Südfall der Ostalpen in 670 m Seehöhe. Die Artbestimmung erfolgte aufgrund der auffällig lila getönten Fellfarbe und der zwischen Kleiner und Großen Hufeisennase liegenden Körpergröße. Diese Beobachtung ist der erste Nachweis dieser Art in Österreich.

Schlüsselwörter

Rhinolophus euryale, Österreich

Lurgrotte Cave (47° 13' N, 15° 22' E; 670 m a.s.l.) is situated 16 km north of Graz. With approximately 6 km length it is the largest cave in the Eastern Alps of Styria. It is located in the Tanneben massif, a 400-500 m thick layer of palaeozoic limestone in the Grazer Bergland. It has two accessible entrances, the upper one in the village of Semriach (Fig. 1) and the lower one in the village of Peggau situated in the Mur valley. A brook crosses the cave with a level difference of 221 m. Lurgrotte Cave is a hibernaculum of many bat species of which the lesser and

The geographic location of the Lurgrotte Cave supports the attribution of the observed specimen to *Rhinolophus euryale*. The distribution range of Mehely's horseshoe bat extends from southern Iberia, south-western France, Sardinia and Sicily to the south-eastern Balkan Peninsula (Bulgaria, Kosovo, and Greece). Few older records are known from coastal Croatia and the borders between Croatia and Bosnia and Herzegovina (RODRIGUES & PALMEIRIM 1999). The western borders of the ranges of Blasius' and Mediterranean horseshoe bat are situated nearer to our observation site (KRYŠTUFEK 1999, IBAÑEZ 1999). The distribution area of Blasius' horseshoe bat extends along the Adriatic coast and covers Albania, Greece, Kosovo and Bulgaria. The Slovenian range of Blasius' horseshoe bat shrank in the 20th century and this species vanished from the northeastern Adriatic coast (KRYŠTUFEK & REŽEK DONEV 2005). *Rh. blasii* is now extinct in the Dinaric Karst of Slovenia (KRYŠTUFEK 2007).

The distribution area of the Mediterranean horseshoe bat comprises the total Mediterranean region and reaches in France and Slovakia locally to more than 45° latitude. In Slovenia, *Rh. euryale* is fairly widespread in the pre-Dinaric region and recent records are reported from the pre-Alpine mountains of the country (KRYŠTUFEK & REŽEK DONEV 2005). KOSELJ (2009) estimated the Slovene population at approximately 1000 individuals. The northernmost Slovenian record is known from Belojača cave in Jelovec pri Makolah, Makole (46 17 N/ 15 39 E) (KOSELJ & AUPUČ 2001) in the Landscape Haloze. The distance between Semriach and the northernmost record in Slovenia is approximately 105 km (Fig. 3). With movements between summer and winter roosts mainly restricted to < 50 km, exceptionally 63-83 km, *Rh. euryale* is thought to be largely sedentary. However, a male was found 134 km from the original site eight years after it has been banded in southern France (GAISLER 2001 a, HUTTERER et al. 2005).

The classification of the sighting of a solitary *Rh. euryale* in Styria/Austria as vagrant, migrant or indication of a northward range expansion is currently not possible. Climate change might have a positive effect on this species with mainly Mediterranean distribution.

Literatur

- DIETZ, C. (2007 a): Blasius Hufeisennase *Rhinolophus blasii* Peters, 1853. Pp. 195-199. In: DIETZ C., von HELVERSEN O. & NILL D. (eds). Handbuch der Fledermäuse Europas und Nordwestafrikas. Franckh Kosmos Verlag, Stuttgart, 399 pp.
- DIETZ, C. (2007 b): Mehely-Hufeisennase *Rhinolophus mehelyi* Matschie, 1901. Pp. 190-194. In: DIETZ, C., von HELVERSEN, O. & NILL, D. (eds): Handbuch der Fledermäuse Europas und Nordwestafrikas. Franckh Kosmos Verlag, 399 pp.
- GAISLER, J. (2001 a): *Rhinolophus euryale* Blasius, 1853 - Mittelmeerhufeisennase. Pp. 59-74. In KRAPP F. (ed): Handbuch der Säugetiere Europas 4., Teil I. Aula Verlag, Wiebelsheim, 602 pp.
- GAISLER, J. (2001 b): *Rhinolophus mehelyi* Matschie, 1901 - Mehely-Hufesennase. Pp.91-104. In: KRAPP F. (ed): Handbuch der Säugetiere Europas 4, Teil I. Aula Verlag, Wiebelsheim, 602 pp.
- HUTTERER, R., IVANOVA, T., MEYER-CORDS, CH. & RODRIGUES, L. (2005): Bat Migrations in Europe. *Naturschutz und Biologische Vielfalt* **28**, 1-162.
- IBAÑEZ, C. (1999): *Rhinolophus euryale* Blasius, 1853. Pp. 92-93. In: MITCHELL-JONES, A. J., AMORI, G., BOGDANOWICZ, W., KRYŠTUFEK, B., REJNDERS, P. H. J., SPITZENBERGER, F., STUBBE, M., THISSEN, J. B. M., VOHRALIK, V. & ZIMA, J. (eds): The Atlas of European Mammals. Poyser Ltd., London, 481 pp.
- KOSELJ, K. & AUPUČ, N. (2001): Prispevek k poznavanju favne netopirjev (Mammalia: Chiroptera) vzhodne Slovenije. *Natura Sloveniae* **3**, 41-62.
- KOSELJ, K. (2009): Južni podkovnjak, *Rhinolophus euryale* Blasius, 1835. Pp 52-54. In: PRESETNIK, P., KOSELJ, K., ZAGMAJSTER, M. (2009). Atlas netopirjev (Chiroptera) Slovenije. Center za kartografijo favne in flore, 151 pp.
- KRYŠTUFEK, B. (1999): *Rhinolophus blasii* Peters, 1866. Pp 90-91. In: MITCHELL-JONES, A. J., AMORI, G., BOGDANOWICZ, W., KRYŠTUFEK B., REJNDERS P. H. J., SPITZENBERGER F., STUBBE M., THISSEN J. B. M., VOHRALIK V. & ZIMA J. (eds): The Atlas of European Mammals. Poyser Ltd., London, 481 pp.
- KRYŠTUFEK, B. (2007): Bat hibernacula in a cave-rich landscape of the north Dinaric karst, Slovenia. *Hystrix, It, J. Mamm. (n. s.)* **18**, 195-204.
- KRYŠTUFEK B. & REŽEK DONEV N. (2005): The Atlas of Slovenian bats (Chiroptera). *Scopolia* **55**, 1-192.
- RODRIGUES, L. & PALMEIRIM, J. (1999): *Rhinolophus mehelyi* Matschie, 1901. Pp. 98-99. In: MITCHELL-JONES, A., AMORI, G., BOGDANOWICZ, W., KRYŠTUFEK, B., REJNDERS, P. H. J., SPITZENBERGER, F., STUBBE, M., THISSEN, J. B. M., VOHRALIK, V. & ZIMA, J. (eds): The Atlas of European Mammals. Poyser Ltd., London, 481 pp.
- SACKL, P., POLT, H., MAYER, A., MOCHÉ, W., MOITZI, F., PASSAUER, U., WEISS, E. & SPITZENBERGER, F. (2006): Artendiversität, Bestandsveränderungen und Quartiermerkmale von Fledermäusen (Mammalia, Chiroptera) in Winterquartieren im Grazer Bergland und in der Grazer Bucht (Steiermark, Österreich). *Mitt. naturwiss. Ver. Steiermark* **141**, 99 – 141.